

# Spikevax<sup>®</sup>

## Messenger ribonucleic acid (mRNA) vaccines

### 1. Moderna Spikevax COVID-19 vaccine available and approved for:

- Primary series (2 doses spaced one month apart)
- Ages 6 months and older
- Booster doses in individuals aged 18 years and older

### 2. Moderna Spikevax Bivalent COVID-19 vaccine (Original/Omicron B.1.1.529 (BA.1)) available and approved for:

- Booster doses
- Ages 6 years and older
- 4 months after the primary series is completed or previous booster dose
- Includes a dose for the original strain and variant BA.1

### 3. Moderna Spikevax Bivalent COVID-19 vaccine (Original/Omicron BA.4/5) available and approved for:

- Booster doses
- Ages 18 years and older
- 4 months after the primary series is completed or previous booster dose
- Includes a dose for the original strain and variant BA.4/5

## Additional information

- Produces a strong immune response
- Proven to be very safe and effective in preventing serious illness and death from COVID-19

## Possible common temporary side effects lasting a few hours to a few days following vaccination:

- redness, soreness, and swelling at the injection site
- chills
- nausea and vomiting
- enlarged lymph nodes
- decreased sense of touch or sensation, numbness, or tingling, itching or pricking sensation
- dizziness
- fatigue
- joint pain
- headache
- mild fever
- muscle aches

# Comirnaty<sup>®</sup>

## Messenger ribonucleic acid (mRNA) vaccines

### 1. Pfizer-BioNTech Comirnaty available and approved for:

- Primary series (2 doses spaced 21 days apart) for ages 6 months and older
- Booster doses (6 months after the primary series) for ages 5 years and older

### 2. Pfizer-BioNTech Comirnaty Original Bivalent (Omicron and BA.1) available and approved for:

- Booster doses
- Ages 5 years and older
- 3-6 months after the primary series is completed or previous booster dose
- Includes a dose for the variant BA.1.

### 3. Pfizer-BioNTech Comirnaty Bivalent (Original and Omicron BA.4/BA.5) available and approved for:

- Booster doses
- Ages 5 years and older
- 3-6 months after the primary series is completed or previous booster dose
- Includes a dose for the variant BA.4 and the variant BA.5.

## Additional information

- Produce a strong immune response
- Proven to be very safe and effective in preventing serious illness and death from COVID-19

## Possible common temporary side effects lasting a few hours to a few days following vaccination:

- redness, soreness, and swelling at the injection site
- chills
- fatigue
- joint pain
- headache
- mild fever
- muscle aches